

# INVESTIGATION ON CHEMICAL CONSTITUENT OF WATER FRACTION OF *PIPER BETLE* L.

## KHẢO SÁT THÀNH PHẦN HOÁ HỌC PHÂN ĐOẠN NƯỚC LOÀI TRẦU KHÔNG

Bui Thi Thu Trang

### ABSTRACT

From water fraction of the whole plant of *Piper betle*, three compounds including thymidine (1), adenosine (2), and 1,2-di-*O*- $\beta$ -glucopyranosyl-4-allylbenzene (3) were isolated. Their chemical structures were elucidated by analyses of their NMR spectral data and comparison with those reported in the literature. Among three isolated compounds, this is the first report the isolation of 1,2-di-*O*- $\beta$ -glucopyranosyl-4-allylbenzene from *P. betle*.

**Keywords:** *Piper betle*, 1,2-di-*O*- $\beta$ -glucopyranosyl-4-allylbenzene, adenosine, thymidine.

### TÓM TẮT

Từ phân đoạn nước của loài trầu không (*Piper betle*) đã được phân lập ba hợp chất gồm thymidine (1), adenosine (2), và 1,2-di-*O*- $\beta$ -glucopyranosyl-4-allylbenzene (3). Cấu trúc hoá học của các hợp chất này được xác định dựa trên phân tích phổ NMR, và so sánh với các số liệu phổ đã được công bố trong tài liệu tham khảo. Trong ba hợp chất phân lập được, đây là lần đầu tiên phân lập được hợp chất 1,2-di-*O*- $\beta$ -glucopyranosyl-4-allylbenzene từ loài trầu không.

**Từ khóa:** Trầu không, 1,2-di-*O*- $\beta$ -glucopyranosyl-4-allylbenzene, adenosine, thymidine.

Faculty of Chemical Technology, Hanoi University of Industry

Email: trangbthoahoc@gmail.com; trangbtt@hau.edu.vn

Received: 12 January 2019

Revised: 07 May 2019

Accepted: 20 February 2020

### 1. INTRODUCTION

*Piper betle* (Piperaceae family) is grown and distributed widely in the tropical countries in the southern and southeastern Asia. The leaves and rhizomes of *P. betle* have been used in traditional medicines for a long time [1]. Chemical components of *P. betle* were reported in the literature including polyphenols, carotenoids, terpenoids, alkaloids, steroids and aliphatic acids [2-6]. The methanolic extract, the essential oil, and several compounds from *P. betle* were screened and showed a lot of valuable bioactivities such as strong anti-microbial and antiseptic, antidotal, anti-inflammatory, and anti-oxidation effects [7-10]. In Vietnam, there were a few investigations focusing on chemical constituents of *P. betle* as well as its essential oil [3-4, 11]. In the contribution to clarify chemical constituent

of *P. betle* growing in Vietnam, herein, we report the isolation and structural determination of three compounds from water fraction of the whole plants of *P. betle*.

### 2. MATERIAL AND METHODS

#### 2.1. General experimental procedures

NMR spectra were recorded on a Jeol 600MHz FT-NMR spectrometer. Column chromatography was performed using a silica gel (Kieselgel 60, 70 - 230 mesh and 230 - 400 mesh, Merck) or RP-18 resins (150 $\mu$ m, YMC), thin layer chromatography (TLC) using a pre-coated silica gel 60 F<sub>254</sub> (0.25mm, Merck) and RP-18 F<sub>254s</sub> plates (0.25mm, Merck).

#### 2.2. Extraction and isolation

The dried and powdered whole plants of *Piper betle* (3kg) was macerated in MeOH for three times (each 6L, 4hrs in ultrasonic bath) to give methanol extract. The methanolic residue (170g) was suspended in water (2L) and successively partitioned with *n*-hexane, dichloromethane, and ethyl acetate. The obtained water soluble fraction was poured on a diaion HP-20 column chromatography (CC), washed with water (2L) and then eluted with gradient solvent system of methanol/water (0 - 100% volume of methanol) to obtain three fractions W1 (6g), W2 (32g) and W3 (17g). The W2 fraction was subjected to a silica gel CC and eluted with gradient solvent system of dichloromethane/ methanol (0 - 100% volume of methanol) to give five fractions W2A-W2F. The W2B was repeatedly chromatographed on a silica gel column, eluting with dichloromethane/methanol/water (6/1/0.1, v/v/v) to give three smaller fractions W2B1-W2B3. Fraction W2B2 was purified on a reverse phase C18 CC using a solvent system of acetone/water (2/3, v/v) to give compound 1 (18 mg) and compound 2 (25mg). The W2D fraction was separated on a silica gel CC with acetone/dichloromethane/water (3/1/0.1, v/v/v) and further purified on a reverse phase C18 CC, eluting with methanol/water (1/1, v/v) to give compound 3 (23mg).

**Thymidine (1):** White amorphous powder; ESI-MS *m/z*: 243 [M+H]<sup>+</sup>; <sup>1</sup>H-NMR (600 MHz, DMSO-*d*<sub>6</sub>)  $\delta$ <sub>H</sub> ppm: 1.86 (3H, s, H-7), 2.21 (2H, m, H-2'), 3.70 (1H, dd, *J* = 3.0, 12.0 Hz, H-5'), 3.77 (1H, dd, *J* = 3.6, 12.0 Hz, H-5'), 3.87 (1H, m, H-4'), 4.37 (1H, m, H-3'), 6.26 (1H, t, *J* = 6.6 Hz, H-1'), 7.79 (1H, s, H-6); <sup>13</sup>C-NMR (150 MHz, DMSO-*d*<sub>6</sub>)  $\delta$ <sub>C</sub> ppm: 12.4 (C-7), 41.2 (C-2'),

62.8 (C-5'), 72.2 (C-3'), 86.2 (C-1'), 88.8 (C-4'), 111.5 (C-5), 138.2 (C-6), 152.4 (C-2), 166.4 (C-4).

**Adenosine (2):** White amorphous powder; ESI-MS  $m/z$ : 268  $[M+H]^+$   $^1H$ -NMR (600 MHz, DMSO- $d_6$ )  $\delta_H$  ppm: 3.55 (1H, m, H-5'), 3.67 (1H, m, H-5'), 3.96 (1H, m, H-4'), 4.12 (1H, dd,  $J = 5.5, 11.0$  Hz, H-3'), 4.61 (1H, dd,  $J = 6.2, 11.0$  Hz, H-2'), 5.87 (1H, d,  $J = 6.2$  Hz, H-1'), 8.13 (1H, s, H-2), 8.35 (1H, s, H-8);  $^{13}C$ -NMR (150 MHz, DMSO- $d_6$ )  $\delta_C$  ppm: 62.2 (C-5'), 71.2 (C-3'), 73.9 (C-2'), 86.4 (C-4'), 88.4 (C-1'), 120.0 (C-5), 140.5 (C-8), 149.7 (C-4), 153.0 (C-2), 156.2 (C-6).

**1,2-di-O- $\beta$ -glucopyranosyl-4-allylbenzene (3):** Pale yellow amorphous powder; ESI-MS  $m/z$ : 497  $[M+Na]^+$   $^1H$ -NMR and  $^{13}C$ -NMR (See Table 1).




Figure 1. Chemical structures of **1-3** from Piper beetle




Figure 2. Important HMBC correlations (H  $\rightarrow$  C) of compound **3**

Table 1. NMR spectral data for compound **3** and reference compound

No.	$^{\#}\delta_C$	$^{a,b}\delta_C$	$^{a,c}\delta_H$ (mult., $J$ in Hz)	No.	$^{\#}\delta_C$	$^{a,b}\delta_C$	$^{a,c}\delta_H$ (mult., $J$ in Hz)	No.	$^{\#}\delta_C$	$^{a,b}\delta_C$	$^{a,c}\delta_H$ (mult., $J$ in Hz)
1	146.8	147.5	-	1-O-Glc			2-O-Glc				
2	148.5	149.2	-	1'	103.3	104.1	4.80 (d, 7.8)	1''	103.5	104.3	4.82 (d, 7.8)
3	120.0	120.9	7.08 (d, 1.8)	2'	74.7	75.1	3.48 (m)	2''	74.7	75.1	3.48 (m)
4	137.3	137.4		3'	77.3	78.2	3.46 (m)	3''	77.3	78.2	3.46 (m)
5	124.7	124.7	6.83 (dd, 1.8; 8.4)	4'	70.9	71.3	3.39 (m)	4''	70.9	71.3	3.39 (m)
6	120.0	120.9	7.15 (d, 8.4)	5'	77.8	77.7	3.32 (m)	5''	77.8	77.7	3.32 (m)
7	40.3	40.6	3.29 (m)	6'	62.0	62.4	3.83 (m) 3.70 (m)	6''	62.0	62.4	3.83 (m) 3.70 (m)
8	138.6	138.8	5.93 (m)								
9	116.3	116.0	5.03 (m)								

Measured in  $^a$ )  $CD_3OD$ ,  $^b$ ) 150MHz,  $^c$ ) 600MHz,  $^{\#}$ ) reported data in  $CD_3OD$  [12]

### 3. RESULTS AND DISCUSSION

Compound **1** was obtained as a white amorphous powder. The  $^1H$ -NMR spectrum of **1** exhibited proton signals at  $\delta_H$  7.79 (1H, s), 6.26 (1H, t,  $J = 6.6$  Hz), 4.37 (1H, m), 3.87 (1H, m), 3.77 (1H, dd,  $J = 3.0, 12.0$  Hz), 3.70 (1H, dd,  $J = 3.6, 12.0$  Hz), 2.21 (2H, m), and methyl group at 1.86 (3H, s). The  $^{13}C$ -NMR spectrum of **1** contained signals of ten carbons at  $\delta_C$  166.4, 152.4, 138.2, 111.5, 88.8, 86.2, 72.2, 62.8, 41.2 and 12.4 ppm. By DEPT spectrum, the signals at  $\delta_C$  62.8 and 41.2 were determined to be  $CH_2$  groups. The chemical structure of **1** was then clearly elucidated by analyses of 2D-NMR spectra such as HSQC and HMBC. Based on the correlations between protons and carbons showed in HSQC spectra, assignments of carbon and proton were obtained including one olefin group at  $\delta_H$  7.79/  $\delta_C$  138.2, one anomeric group at  $\delta_H$  6.26/  $\delta_C$  86.2, two oxymethine groups at  $\delta_H$  4.37/  $\delta_C$  72.2 and  $\delta_H$  3.87/  $\delta_C$  88.8, one oxymethylene group at  $\delta_H$  3.77 and 3.70/  $\delta_C$  62.8, one methylene group at  $\delta_H$  2.21/  $\delta_C$  41.2, and one methyl group at  $\delta_H$  1.86/  $\delta_C$  12.4. Three remaining carbon signals at  $\delta_C$  166.4, 152.4, and 111.5 did not observe correlations with proton in the HSQC spectrum which were assigned for nonprotonated carbons. An anomeric carbon signal at  $\delta_C$  86.2 and HMBC correlation between anomeric proton ( $\delta_H$  6.26) and carbon C-2 ( $\delta_C$  152.4), C-6 ( $\delta_C$  138.2) suggested for the N-glycoside linkage. The HMBC correlation between methyl protons ( $\delta_H$  1.86) and C-6 ( $\delta_C$  138.2)/ C-5 ( $\delta_C$  111.5)/ C-4 ( $\delta_C$  166.4) indicated the assignment of C-4/C-5/C-6 and a double bond at C-5/C-6. In addition, the chemical shift of C-4 ( $\delta_C$  166.4) allowed to determine carbonyl group of C-4. Also, chemical shift of C-2 at  $\delta_C$  152.4 ppm can be predicted other carbonyl group at C-2 and the linkage between C-2 and C-4 via a nitrogen atom. Five carbon signals of monosaccharide suggested the presence of a pentose. In addition, signals of methylene group C-2' ( $\delta_H$  2.21/  $\delta_C$  41.2) suggested for the presence of a 2-deoxypentose. The NMR data of **1** was compared and

observed in good agreement with those reported of thymidine in the literature [13]. Consequently, compound **1** was determined to be thymidine.

Compound **2** was isolated as a white amorphous powder. The  $^1\text{H-NMR}$  spectrum of **2** exhibited the signals at  $\delta_{\text{H}}$  8.13 (s), 8.35 (s), 5.87 (d,  $J = 6.2$  Hz), 4.61 (dd,  $J = 6.2, 11.0$  Hz), 4.12 (dd,  $J = 5.5, 11.0$  Hz), 3.96 (m), 3.67 (m), and 3.55 (m). The  $^{13}\text{C-NMR}$  spectrum of **2** showed ten carbons at  $\delta_{\text{C}}$  156.2, 153.0, 149.7, 140.5, 120.0, 88.4, 86.4, 73.9, 71.2, 62.2. By DEPT spectrum, these signals were separated into one  $\text{CH}_2$  group at  $\delta_{\text{C}}$  62.2, three non-protonated carbons at  $\delta_{\text{C}}$  156.2, 149.7, 120.0; and remaining six CH groups. Among them, five carbon signals at  $\delta_{\text{C}}$  88.4, 73.9, 71.2, 86.4, 62.2 suggested for the presence of a ribose moiety. The chemical shift of C-1' ( $\delta_{\text{C}}$  88.4), and HMBC correlation from H-1' ( $\delta_{\text{H}}$  5.87) to C-4 ( $\delta_{\text{C}}$  149.7), C-8 ( $\delta_{\text{C}}$  140.5) suggested the presence of a N-glycoside linkage. In the aglycone moiety, there were only two signals of  $\text{sp}^2$  methine groups at  $\delta_{\text{H}}$  8.13/  $\delta_{\text{C}}$  153.0 and  $\delta_{\text{H}}$  8.35/  $\delta_{\text{C}}$  140.5 which were suggested for the structure of adenine alkaloid. The  $^{13}\text{C-NMR}$  spectral data of **2** was also in good consistence with that reported data of adenosine in the literature [14]. The NMR assignment of **2** was confirmed again by HMBC spectrum. Example, HMBC correlations between anomeric proton H-1' ( $\delta_{\text{H}}$  5.87) and C-4 ( $\delta_{\text{C}}$  149.7), C-8 ( $\delta_{\text{C}}$  140.5) indicated the assignment of chemical shift values of C-4 and C-8, respectively. HMBC correlations between H-8 ( $\delta_{\text{H}}$  8.35) and C-4 ( $\delta_{\text{C}}$  149.7)/ C-5 ( $\delta_{\text{C}}$  120.0) were demonstrated for assignment of C-5. And HMBC correlations between H-2 ( $\delta_{\text{H}}$  8.13) and C-4 ( $\delta_{\text{C}}$  149.7)/ C-6 ( $\delta_{\text{C}}$  156.2) were supported for assignment of C-6. Similarly, the signals of monosaccharide moiety were assigned by 2D-spectra and spin-spin coupling constant observing in  $^1\text{H-NMR}$  spectrum. Oxymethylene group was deduced by the signal at  $\delta_{\text{H}}$  3.67, 3.55/  $\delta_{\text{C}}$  62.2. HMBC correlations from H-5' ( $\delta_{\text{H}}$  3.67, 3.55) to C-4' ( $\delta_{\text{C}}$  86.4), C-3' ( $\delta_{\text{C}}$  71.2) consisted to the data of C-3' and C-4'. Then, the remaining oxymethine signal at  $\delta_{\text{C}}$  73.9 was assigned to C-2'. Adenosine was one of nucleoside. This compound was known as a main component of *Cordyceps sinensis*, played an important role in circulatory system, controlling immune system [15].

Compound **3** was isolated as a pale yellow amorphous powder. The  $^1\text{H-NMR}$  spectrum of **3** exhibited three aromatic protons of ABX spin coupled system at  $\delta_{\text{H}}$  7.08 (1H, d,  $J = 1.8$  Hz), 6.83 (1H, dd,  $J = 1.8; 8.4$  Hz), and 7.15 (1H, d,  $J = 8.4$  Hz), indicating the presence of a 1,3,4-three substituted phenyl group. The vinyl ( $\text{CH}_2=\text{CH}$ ) group were recognized by three olefinic proton signals at  $\delta_{\text{H}}$  5.93 (1H, m) and 5.03 (2H, m). Two anomeric protons signals were showed at  $\delta_{\text{H}}$  4.80 and 4.82 (each 1H, d,  $J = 7.8$  Hz) suggested the presence of two monosaccharide units. Analysis of  $^{13}\text{C-NMR}$  spectrum of **3** revealed twenty-one carbons. Among them, twelve carbons at  $\delta_{\text{C}}$  104.1/ 104.3, 75.1/ 75.1, 78.2/ 78.2, 71.3/ 71.3, 77.7/ 77.7, and 62.4/ 62.4 characterised for two glucopyranosyl groups. At the same

time, the spin-spin coupling constant of anomeric proton ( $J = 7.8$  Hz) indicated the  $\beta$ -glucopyranosyl linkages. Next, HMBC correlations between H-7 ( $\delta_{\text{H}}$  3.29) and C-3 ( $\delta_{\text{C}}$  120.9)/ C-4 ( $\delta_{\text{C}}$  137.4)/ C-5 ( $\delta_{\text{C}}$  124.7)/ C-8 ( $\delta_{\text{C}}$  138.8)/ C-9 ( $\delta_{\text{C}}$  116.0) indicated the location of propenyl group at C-4. In addition, HMBC correlations between H-3 ( $\delta_{\text{H}}$  7.08)/ H-5 ( $\delta_{\text{H}}$  6.83) and C-1 ( $\delta_{\text{C}}$  147.5) and downfield chemical shift of C-1 ( $\delta_{\text{C}}$  147.5) indicated an oxygenated carbon C-1. Similarly, HMBC correlations between proton H-6 ( $\delta_{\text{H}}$  7.15) and C-4 ( $\delta_{\text{C}}$  137.4)/ C-2 ( $\delta_{\text{C}}$  149.2), in combination with downfield chemical shift of C-2 ( $\delta_{\text{C}}$  149.2) suggested another oxygenated  $\text{sp}^2$  carbon C-2. Finally, HMBC correlations between anomeric protons H-1' ( $\delta_{\text{H}}$  4.80) and C-1 ( $\delta_{\text{C}}$  147.5), H-1'' ( $\delta_{\text{H}}$  4.82) and C-2 ( $\delta_{\text{C}}$  149.2) were confirmed the positions of two glucopyranosyl groups at C-1 and C-2. Consequently, compound **3** was determined to be 1,2-di-O- $\beta$ -glucopyranosyl-4-allylbenzene. The  $^{13}\text{C-NMR}$  data of **3** was well consisted with those reported data of 1,2-di-O- $\beta$ -D-glucopyranosyl-4-allylbenzene in the literature (Table 1). This compound was isolated previously from *Alpinia officinarum*, a species belonging Zingiberaceae family and was expected causing anti-oxidation activity of *Alpinia officinarum* [12]. However, until now, this is the first report on the isolation of 1,2-di-O- $\beta$ -glucopyranosyl-4-allylbenzene from *Piper betle*.

#### 4. CONCLUSIONS

Three compounds including thymidine, adenosine, and 1,2-di-O- $\beta$ -glucopyranosyl-4-allylbenzene were isolated from water soluble fraction of the whole plants of *P. betle*. Their NMR spectral data were well consisted with those reported in the literature. To the best of our knowledge, this is the first report on the isolation of 1,2-di-O- $\beta$ -glucopyranosyl-4-allylbenzene from *P. betle*.

#### REFERENCE

- [1]. Lợi Đ.T., 2003. *Những cây thuốc và vị thuốc Việt Nam*. NXB Y học, Hà Nội,
- [2]. Panuwat S, Natcha SE, Panchuti P., 2006. *Antimicrobial and antioxidant activities of essential oil of Betel (Piper betel Linn.)*. Natural Science, 91-100.
- [3]. Thanh L, Dũng NX, Piet AL., 1996. *Thành phần hóa học của tinh dầu lá trầu không (Piper betle L)*. Tạp chí Dược liệu, 18-20.
- [4]. Chính PT., 2009. *Nghiên cứu thành phần hóa học và hoạt chất sinh học của lá trầu (Piper betle L) miền bắc Việt Nam*. Luận văn thạc sỹ khoa học, Trường Đại học khoa học Tự nhiên, Đại học Quốc gia Hà Nội.
- [5]. Ghosh K, Bhattacharya TK., 2005. *Chemical constituents of Piper betle Linn. (Piperaceae) roots*. Molecules, 10, 798-802.
- [6]. Lim CM, Ee G, Rahmani M, Bong CFJ., 2009. *Alkaloids from Piper nigrum and Piper betle*. Pertanika J. Sci. & Technol., 17, 149-154.
- [7]. Thuraijajah N, Rahim Z., 2006. *Effect of Piper betle L. Leaf extract on the virulence activity of Streptococcus mutans - An in vitro Study*. Pakistan J. Biol. Sci., 9, 1470-1475.

[8]. Nalina T, Rahim ZHA., 2007. *The crude aqueous extract of Piper betle L. and its antibacterial effect towards Streptococcus mutans*. American J Biotechnol. Biochem., 3, 10-15.

[9]. Bhattacharya S, Banerjee D, Bauri AK, Chattopadhyay S, Bandyopadhyay SK., 2007. *Healing property of the Piper betel phenol, allylpyrocatechol against indomethacin-induced stomach ulceration and mechanism of action*. World J Gastroenterol, 13, 3705-3713.

[10]. Zeng HW, Jiang YY, Cai DG, Bian J, Long K, Chen ZL., 1997. *Piperbetol, methylpiperbetol, piperol A and piperol B: a new series of highly specific PAF receptor antagonists from Piper betle*. Planta Med., 63, 296-298.

[11]. Thẩm PT, Yến LTH, Hùng TT, Thanh NT, Chính PT., 2013. *Phân lập các hợp chất trong thân cây trầu không (Piper betle L.)*. Tạp chí Khoa học và Công nghệ, Đại học Thái Nguyên, 112, 27-33.

[12]. Ly TN, Yamauchi R, Shimoyamada M, Kato K., 2002. *Isolation and structural elucidation of some glycosides from the rhizomes of smaller galanga (Alpinia officinarum Hance)*. J. Agric. Food Chem., 50, 4919-4924.

[13]. Moyroud E, Biala E, Strazewski P., 2000. *Synthesis and enzymatic digestion of an RNA nonamer in both enantiomeric forms*. Tetrahedron, 56, 1475-1484.

[14]. Song MC, Yang HJ, Jeong TS, Kim KT, Baek NI., 2008. *Heterocyclic compounds from Chrysanthemum coronarium L. and their inhibitory activity on hACAT-1, hACAT-2, and LDL-oxidation*. Arch Pharm Res, 31, 573-578.

[15]. Tsai YJ, Lin LC, Tsai TH., 2010. *Pharmacokinetics of adenosine and cordycepin, a bioactive constituent of Cordyceps sinensis in rat*. J. Agric. Food Chem., 58, 4638-4643.

---

#### THÔNG TIN TÁC GIẢ

**Bùi Thị Thu Trang**

Khoa Công nghệ Hóa, Trường Đại học Công nghiệp Hà Nội